

TRIPLO



General production advice ware potatoes

- Suitable to make (home) fries
- Multi purpose variety
- Good Blight resistance



Morphology

| | |
|---------------------|--------------------|
| Cooking type | B - Slightly mealy |
| Maturity | 70 Medium early |
| Yield mature | 104 Moderate high |
| Tuber size | 83 Large |
| Tuber shape | Round oval / Oval |
| Number of tubers | 9-11 |
| Flesh after cooking | Light yellow |
| Skin colour | Yellow |
| Berries | Few berries |

Tolerances

| | |
|------------------------|--------------------|
| Dormancy period | 54 Moderate short |
| Emergence | 80 Fast |
| Metribuzin sensitivity | 51 Sensitive |
| Foliage development | 75 Strong |
| Internal bruising | 8 Little sensitive |
| Little Potato disorder | 80 Not sensitive |
| Dry matter content | 20,2 |
| UWW | 385 |

Resistances

| | |
|----------------------|----------------------|
| Spraing | Slightly susceptible |
| Foliage Blight | Slightly susceptible |
| Tuber Blight | Resistant |
| Common scab | Susceptible |
| Powdery scab | Very susceptible |
| PVYn | Slightly susceptible |
| Yntn tuber tolerance | Slightly sensitive |

PCN Resistance

| Ro1 | Ro2/3 | Pa2 | Pa3 |
|--------------|-------|-----|-----|
| 9 | 3 | | |
| Wart disease | | | |
| F1 | F2 | F6 | F18 |
| 10 | 3 | 3 | 3 |

Fertilizer

- Nitrogen (N): approx. 200 kg N/ha inclusive of soil supply.
- Potassium and phosphate fertilisation as standard advice.

TRIPLO



General production advice ware potatoes

Pre-treatment and planting

- TRIPLO has a short dormancy.
- If an apical sprout is present, remove it and plant with small white buds.
- Pre-sprouting is not necessary.
- Avoid soils with high common scab pressure.
- Plant large seed deeper to avoid greening.
- Do not plant in cold soils in relation to the risk of irregular emergence.

Plant populations

| Seed size | Plant population/ha | Row distance | |
|-----------|---------------------|--------------|-------|
| | | 75 cm | 90 cm |
| -28/35 | 60.000 | 22 | 19 |
| -35/50 | 50.000 | 27 | 22 |
| -35/55 | 48.000 | 28 | 23 |
| -50/60 | 34.000 | 39 | 33 |

-Plant depth: 3 cm deeper than normal.



Growing attentionpoints

- TRIPLO grows high in the ridge
- Variety is quite sensitive to metribuzin (Sencor), only use it pre-emergence.
- With risk of Common Scab, irrigate the crop at tuber initiation.

Haulmkillling and harvest

- TRIPLO is quite early maturing, haulmkillling shouldn't give any problems.

Storage

- Dry quick and keep it dry to avoid Silver scurf
- TRIPLO is little susceptible to Fusarium.