

SPUNTA



Production advice ware potatoes NORTH AFRICA



Morphology

Cooking type	B - Slightly mealy
Maturity	64 Medium late
Yield mature	114 High
Tuber size	85 Large
Tuber shape	Long oval
Number of tubers	9-11
Flesh after cooking	Light yellow
Skin colour	Yellow
Berries	No berries

Tolerances

Dormancy period	62 Medium
Emergence	80 Fast
Metribuzin sensitivity	75 Moderately sensitive
Foliage development	79 Very strong
Internal bruising	14 Quite sensitive
Little Potato disorder	50 Quite Sensitive
Dry matter content	20,0
UWW	366

Resistances

Spraing	Highly resistant
Foliage Blight	Susceptible
Tuber Blight	Susceptible
Common scab	Susceptible
Powdery scab	Very susceptible
PVYn	Susceptible
Yntn tuber tolerance	Slightly sensitive

PCN Resistance

Ro1	Ro2/3	Pa2	Pa3
3	1	2	2

Wart disease

F1	F2	F6	F18
10			

Fertilizer

SPUNTA



Production advice ware potatoes NORTH AFRICA

Pre-treatment and planting

Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm



Growing attentionpoints

Haulmkillling and harvest

Storage